## **CHAPTER 10 SOLUTION SHEET-20151110**

## Hypothesis Testing for Two Means, Two proportions, Matched Pairs

What type of test? (circle one)	SIEP 3: Test Statistic: indicate t or z; state value: t or z (circle one) =
Two proportions Two means, independent samples Two means, paired samples	p-value =
What are the population parameters? (circle one) $P_1 \& P_2 \qquad \mu_1 \& \mu_2 \qquad \mu_d$	Use the previous information to draw the graph of this situation.
What is the random variable? (circle one) $P'_1 - P'_2 = \overline{X}_1 - \overline{X}_2 = \overline{X}_d$ In words for this problem's scenario, CLEARLY describe the population parameter(s):	Label & scale both horizontal axes (for the random variable and for the test statistic t or z)
	Shade and label the region(s) corresponding to the p-value
STEP 1: Ho: Ha:	STEP 4:
Significance Level $\alpha =$	Indicate the decision ("reject Ho" or "do not reject Ho"), and the reason
STEP 2: Distribution to use for the test:	decision reason for decision
Normal N(,)  OR t with df =	
Calculator Test to use: 2 SamT Test 2 SamZTest TTest 2PropZTest	
In complete sentences, explain how you determined which distribution to use.	
	IS THE RESULT STATISTICALLY SIGNIFICANT?
	STEP 5: Write an appropriate conclusion, in the context of the problem
Values of the Sample Statistics as appropriate for this test. $\overline{x_d} = \underline{\hspace{1cm}}$	using COMPLETE SENTENCES, stating the significance level and whether or not the sample data contain sufficient evidence to support the alternate hypothesis. Conclusion:
$OR \ \overline{x}_1 - \overline{x}_2 = \underline{\qquad} - \underline{\qquad} = \underline{\qquad}$	
with $s_1 = $ $n_1 = $ $s_2 = $ $n_2 = $	
<b>OR</b> p' <sub>1</sub> - p' <sub>2</sub> ===	
with $x_1 = \underline{\hspace{1cm}} n_1 = \underline{\hspace{1cm}} x_2 = \underline{\hspace{1cm}} n_2 = \underline{\hspace{1cm}}$	